

#### **ARSET**

Applied Remote Sensing Training http://arset.gsfc.nasa.gov

# Creating and Using Normalized Difference Vegetation Index (NDVI) from Satellite Imagery

Instructors: Cindy Schmidt and Amber McCullum

Week 3

### Course Structure

- One lecture per week every Wednesday from February 10 to March 2 at 12:00-1:00pm EST (-05:00 UTC)
  - Lectures
  - In-class exercise
  - Q&A
  - Homework exercises
- Webinar recordings, PowerPoint presentations, in-class exercises, and homework assignments can be found after each session at:
  - http://arset.gsfc.nasa.gov/ecoforecasting/webinars/advanced-webinar-creating-and-usingnormalized-difference-vegetation-index
- Q&A: Following each lecture and/or by email (<u>cynthia.l.schmidt@nasa.gov</u>)

### Homework and Certificates


- Homework
  - Hands-on exercise each week
  - Answers must be submitted via Google Form
- Certificate of Completion:
  - Attend all 4 webinars
  - Complete all 4 homework assignments by the deadline (access from ARSET website)
    - Week 3 Deadline: Wednesday March 9th
  - You will receive certificates approximately 2 months after the completion of the course from: <a href="mailto:martins@ssaihq.com">martins@ssaihq.com</a>



### **Accessing Course Materials**

<a href="http://arset.gsfc.nasa.gov/ecoforecasting/webinars/advanced-webinar-creating-and-using-normalized-difference-vegetation-index">http://arset.gsfc.nasa.gov/ecoforecasting/webinars/advanced-webinar-creating-and-using-normalized-difference-vegetation-index</a>







Course materials are provided here using each specified link and will be active after each week


### **Course Outline**


### Week 1

Overview of NDVI and **QGIS** 









### Week 2

**NDVI** with Landsat

### Week 4

**MODIS** NDVI **Anomaly Mapping** 




Week 3

**NDVI** Time **Series** 



### Week 3 Agenda

- Overview of MODIS Imagery
- Live Demos: Global Agricultural Monitoring (GLAM) Project
  - MODIS/NDVI Time Series
     Database
  - GIMMS MODIS system
- Q&A




MODIS NDVI of Nile River Basin from GLAM

### **NDVI** Time Series and Anomalies


- Can be used to identify
  - Changes in vegetation health
  - Land degradation
  - Decreased/increased precipitation
  - Changes in phenology (green-up timing)

NDVI anomaly, August 1984, Ethiopia



NDVI time series for 2015, Ethiopia







NDVI anomaly, August 2004, USA




### **MODIS Characteristics**

- Spatial Resolution
  - 250m, 500m, 1km
- Temporal Resolution
  - Daily, 8-day, 16-day, monthly, quarterly, yearly
  - 2000-present
- Data Format
  - Hierarchal data format









- Spectral Coverage
  - 36 bands (major bands include Red, Blue, IR, NIR, MIR)
    - Bands 1-2: 250m
    - Bands 3-7: 500m
    - Bands 8-36: 1000m
- Orbital gaps

# MODIS Tiles vs. Landsat



### **MODIS Naming Convention**

MODIS filenames follow a naming convention which gives useful information regarding the specific product. For Example:



\*\*NOTE: MOD – Terra; MYD – Aqua; MCD - Combined

# **MODIS Land Products**

| MODIS<br>Name | Product Name Short name   | Spatial Resolution (m) | Temporal        |
|---------------|---|------------------------|-----------------|
| MOD 09        | Surface Reflectance   | 500                    | 8-day           |
| MOD 11        | Land Surface Temperature  | 1000                   | Daily, 8-day    |
| MOD 12        | Land Cover/Change   | 500                    | 8-day, Yearly   |
| MOD 13        | Vegetation Indices  | 250-1000               | 16 day, monthly |
| MOD 14        | Thermal Anomalies/Fire  | 1000                   | Daily, 8-day    |
| MOD 15        | Leaf Area Index/Fraction of Absorbed Photosynthetically Active Radiation (FPAR) | 1000                   | 4-day, 8-day    |
| MOD 16        | Evapotranspiration  |                        |                 |
| MOD 17        | Primary Production  | 1000                   | 8-day, yearly   |
| MOD 43        | Bidirectional reflectance distribution function (BRDF)/Albedo                   | 500-1000               | 16-day          |
| MOD 44        | Vegetation Continuous Fields  | 250                    | yearly          |
| MOD 45        | Burned Area   | 500                    | monthly         |

#### All MODIS Land Products are available at processing Level 3


### Where to Obtain MODIS Land Products

 Land Processes Distributed Active Archive (LP DAAC)

> https://lpdaac.usgs.gov/ dataset\_discovery/modis/ modis products table

Earth Observing System
 Data and Information
 System (EOSDIS):

http://Earthdata.nasa.gov



### Where to Obtain MODIS Land Products



ECHO Reverb: http://reverb.echo.nasa.gov



Data Subsetting and Visualization: Oakridge National Lab DAAC (ORNL DAAC): <a href="http://daac.ornl.gov">http://daac.ornl.gov</a>



GLCF: http://www.landcover.org/data/lc



GLOVIS: <a href="http://glovis.usgs.gov">http://glovis.usgs.gov</a>



Fire Information for Resource Management System (FIRMS): <a href="https://earthdata.nasa.gov/data/near-real-time-data/firms">https://earthdata.nasa.gov/data/near-real-time-data/firms</a>

### Where to Obtain MODIS Land Products



Worldview (Fires, Land Surface Temperature and Snow Cover): <a href="https://earthdata.nasa.gov/labs/worldview">https://earthdata.nasa.gov/labs/worldview</a>



Visualization, SERVIR:


https://www.servirglobal.net/Global/MapsData/ InteractiveMapper.aspx



MRTWeb: <a href="http://mrtweb.cr.usgs.gov">http://mrtweb.cr.usgs.gov</a>

### **MRTWeb**

- MRTWeb enables you to:
  - Visualize
  - Select
  - Mosaic
  - Subset
  - Reproject
  - Reformat



## What is EVI?

- Enhanced Vegetation Index (MOD13Q1) standard product from MODIS
  - More sensitive to changes in areas having high biomass
  - Reduces the influence of atmospheric conditions on vegetation index values
  - Corrects for canopy background signals




Image credit: NASA Earth Observatory


### What is EVI?

EVI is calculated as:

$$EVI = G * (NIR-RED)/NIR+C1*RED-C2*BLUE+L$$

- Red and Blue bands must be atmospherically corrected
- G is a scaling factor. G = 2.5
- C<sub>1</sub>, C<sub>2</sub> and L are coefficients to correct for atmospheric condition. For MODIS and Landsat C<sub>1</sub>=6, C<sub>2</sub> = 7.5 and L = 1

Comparison between MODIS NDVI and EVI in North America
March 2000



Source: earthobservatory.nasa.gov



### Global Agriculture Monitoring (GLAM) Project

- Global 16-day 250 m NDVI time series database
- USDA and NASA collaboration
  - Integrates remote sensing into USDA monitoring system
- Web-interface
  - Plot time series graphs over crop season
  - Monitor current conditions
  - Spatially view NDVI anomalies
  - Plot histograms of current and historical NDVI data




#### Website:

http://pekko.geog.umd.edu/usda/test/

## Global Agriculture Monitoring (GLAM) Project

- Global Inventory Modeling and Mapping Studies (GIMMS) MODIS monitoring system
- USDA and NASA collaboration
  - Integrates remote sensing into USDA monitoring system
- Web-interface
  - Global NDVI distribution datasets in online viewer
  - Download GeoTIFF imagery
  - Spatially view NDVI and NDVI anomalies



#### Website:

http://glam1.gsfc.nasa.gov

### Contacts

- ARSET Land Management and Wildfire Contacts
  - Cynthia Schmidt: Cynthia.L.Schmidt@nasa.gov
  - Amber McCullum: AmberJean.Mccullum@nasa.gov
- General ARSET Inquiries
  - Ana Prados: aprados@umbc.edu
- ARSET Website:
  - http://arset.gsfc.nasa.gov/



#### **ARSET**

Applied Remote Sensing Training http://arset.gsfc.nasa.gov

# **Thank You**

**Next Week:** 

**MODIS NDVI Anomalies**